

Dear Author,

Please, note that changes made to the HTML content will be added to the article before publication, but are not reflected in this PDF.


Note also that this file should not be used for submitting corrections.



Contents lists available at ScienceDirect

Computers & Fluids

journal homepage: www.elsevier.com/locate/complfluid



Lie symmetry analysis, nonlinear self-adjointness and conservation laws to an extended (2+1)-dimensional Zakharov–Kuznetsov–Burgers equation [☆]

Gangwei Wang ^{a,*}, K. Fakhar ^b

^a School of Mathematics and Statistics, Beijing Institute of Technology, Beijing 100081, PR China

^b Department of Mathematics, University of British Columbia, Vancouver V6T 1Z2, Canada

ARTICLE INFO

Article history:

Received 6 July 2014

Received in revised form 28 May 2015

Accepted 26 June 2015

Available online xxxxx

Keywords:

(2+1)-dimensional Zakharov–Kuznetsov–

Burgers equation

Lie symmetry analysis

Nonlinear self-adjointness

Conservation laws

ABSTRACT

This paper addresses an extended (2+1)-dimensional Zakharov–Kuznetsov–Burgers (ZKB) equation. The Lie symmetry analysis leads to many plethora of solutions to the equation. The nonlinear self-adjointness condition for the ZKB equation is established and subsequently used to construct simplified but infinitely many nontrivial and independent conserved vectors. In particular, we also get conservation laws of the equation with the corresponding Lie symmetry.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

The nonlinear evolution equations (NLEEs) are encountered in a variety of scientific fields, such as physics, chemistry, engineering and others. A vast amount of research work has been investigated in the study of exact solutions of the NLEEs, in particular, the solitary wave solutions largely due to their frequent occurrence in nature. Besides their physical relevances, they serve as a bench mark for accuracy of numerical schemes as well as prove to be quiet handy in testing of computer algorithms. Due to the increased interest in the NLEEs, a broad range of analytical and numerical methods have been developed to construct exact solutions to NLEEs. Some of these efficient methods are the Lie symmetry method [1–3], Darboux transformation method [4], Jacobi elliptic method [5], Painleve analysis [6], the inverse scattering method [7], the Baklund transformation method [8], the conservation law method [9], the Hirota bilinear method [10], the ansatz method [11] and many other methods.

Zakharov and Kuznetsov [12] established an equation which is related to nonlinear ion-acoustic waves (IAWs) in magnetized

plasma including cold ions and hot isothermal electrons. Few related studies concerning Zakharov and Kuznetsov and its generalized form are discussed in [13–19]. The quantum hydrodynamic (QHD) model is a generalization of the classical fluid model of plasmas where QHD transport equations are expressed with reference to conservation laws for particle momentum and energy. Several authors for instance, Stenflo et al. [20], Khan et al. [21] etc., have used QHD model to study the linear and nonlinear waves in quantum plasma. Further, the existence of a small number of ions along with the electrons and positron in many astrophysical environments, attracted much attention and consequently lot of work available in the literature (e.g., [22,23]). It is well know that the small amplitude wave propagation in a medium which posses both the characteristics dispersive and dissipative terms can be best modeled by Korteweg–deVries–Burgers (KdVB) equation. Mamun and Shukla [24] and Shukla and Mamun [25] in their study observed that the dissipative Burger term in KdVB was due to the presence of kinematic viscosity in the plasma. Also, they were able to generate dispersive shock wave in the plasma. El-Bedwehy and Moslem derived the Zakharov Kuznetsov Burgers (ZKB) equation [26] in an electron–positron–ion (e–p–i) plasma and applied their numerical results to the electrostatic fluctuations in the interstellar medium. Masood et al. [27] studies the obliquely propagating nonlinear quantum ion acoustic shock wave in a viscous quantum (e–p–i) magnetoplasma. They have used QHD model with the small amplitude expansion method to independently

[☆] The project is supported by the Research Project of China Scholarship Council (No. 201406030057), National Natural Science Foundation of China (NNSFC) (Grant No. 11171022), Graduate Student Science and Technology Innovation Activities of Beijing Institute of Technology (No. 2014cx10037).

* Corresponding author.

E-mail address: pukai1121@163.com (G. Wang).

derive the ZKB equation. They have used tanh method to obtain the results.

In this paper, we consider the (2+1)-dimensional ZKB equation and investigate it from the point view of Lie symmetries and conservation laws. By omitting the details of derivation, we directly write ZKB equation in the form

$$u_t + auu_x + bu_{xxx} + cu_{xyy} - du_{xx} - eu_{yy} = 0, \quad (1)$$

where a, b, c, d and e , with d and e being positive, are constant quantities which involves the physical quantities like: mass; density; magnetic field; kinematic viscosity; plasma frequency; superthermality; ion gyrofrequency, etc., while x, y and t are the independent variables that represent the spatial and temporal variables respectively, where as $u(x, y, t)$ is the dependent variable that represents the wave profile. For detail discussions reader is refer to [26,27].

The paper is organized in the following manner. In Section 2, similarity reductions and explicit solutions are derived. In Section 3, we will show that this equation is nonlinearly self-adjoint. On the basis of the point symmetries, conservation laws are constructed. Finally, the main findings of the paper are recapitulated in Section 4.

2. Similarity reductions and exact solutions

The corresponding vector fields of (1) are

$$V_1 = \frac{\partial}{\partial x}, \quad V_2 = \frac{\partial}{\partial y}, \quad V_3 = \frac{\partial}{\partial t}, \quad V_4 = t \frac{\partial}{\partial x} + \frac{1}{a} \frac{\partial}{\partial u}. \quad (2)$$

We make some discussions on the ZKB equation based on the vector fields.

(I) V_1

For the generator V_1 , we have

$$u = f(g, h), \quad (3)$$

where $g = y, h = t$ are the group-invariants. Substituting (3) into (1), one can get

$$f_h - ef_{gg} = 0. \quad (4)$$

It is important to note that (4) is the celebrated Heat equation.

(II) V_2

For the generator V_2 , we get

$$u = f(r, h), \quad (5)$$

where $r = x, h = t$ are the group-invariants. Substituting (5) into (1), we reduce it to the following PDE

$$f_h + aff_r + bf_{rr} - df_{rr} = 0. \quad (6)$$

(III) V_3

For the generator V_3 , we get

$$u = f(r, g), \quad (7)$$

where $r = x, g = y$ are the group-invariants. Putting (7) into (1), one can obtain

$$aff_r + bf_{rr} + cf_{rgg} - df_{rr} - ef_{gg} = 0. \quad (8)$$

(IV) V_4

In this case, one can obtain

$$u = f(h, g) + \frac{x}{at}, \quad (9)$$

where $h = t, g = y$ are the group-invariants. Substituting (9) into (1), one can obtain

$$hf_h + f - ehf_{gg} = 0. \quad (10)$$

(V) $V_1 + V_2$

The symmetry $V_1 + V_2$ yields the following invariants

$$h = t, \quad s = x - y, \quad u = f(h, s). \quad (11)$$

Treating f as the new dependent variable and h and s as new independent variables, the ZKB Eq. (1) transforms to

$$f_h + aff_s + (b + c)f_{sss} - (d + e)f_{ss} = 0. \quad (12)$$

It is should be noted that all above reduced equation are (1+1)-dimensional PDEs. It is also difficult to get solutions of them. In order to get their solutions, once again, by using Lie symmetry method. For the sake of simplicity, in what follows, we only consider (12) in details.

The symmetry algebra of (12) is generated by the vector fields

$$Y_1 = \frac{\partial}{\partial h}, \quad Y_2 = \frac{\partial}{\partial s}, \quad Y_3 = h \frac{\partial}{\partial s} + \frac{1}{a} \frac{\partial}{\partial f}. \quad (13)$$

The combination $Y_1 + \lambda Y_2$ of the two symmetries Y_1 and Y_2 yields the following invariants

$$f = \Psi(\theta), \quad \theta = s - \lambda h, \quad (14)$$

and using (14) and (12) is transformed to the nonlinear ODE

$$-\lambda \Psi'' + a \Psi \Psi' + (b + c) \Psi''' - (d + e) \Psi'' = 0. \quad (15)$$

Now, we search a solution of (15) in a power series of the form [28,29]

$$\Psi = \sum_{n=0}^{\infty} c_n \theta^n. \quad (16)$$

Substituting (16) into (15), one can get

$$\begin{aligned} & -\lambda c_1 - \lambda \sum_{n=1}^{\infty} (n+1) c_{n+1} \theta^n + ac_0 c_1 + a \sum_{n=1}^{\infty} \sum_{j=0}^n (n+1-j) c_j c_{n+1-j} \theta^n \\ & + 2(d+e)c_2 + (d+e) \sum_{n=1}^{\infty} (n+1)(n+2) c_{n+2} \theta^n + 6(b+c)c_3 \\ & + (b+c) \sum_{n=1}^{\infty} (n+3)(n+2)(n+1) c_{n+3} \theta^n = 0. \end{aligned} \quad (17)$$

Next, from (17), for the case of $n = 0$, one gets

$$c_3 = \frac{\lambda c_1 - ac_0 c_1 - 2(d+e)c_2}{6(b+c)}. \quad (18)$$

Generally, for $n \geq 1$, one obtains

$$\begin{aligned} c_{n+3} = & \frac{1}{(b+c)(n+1)(n+2)(n+3)} \left(\lambda(n+1)c_{n+1} - (d+e)(n+1)(n+2) \right. \\ & \left. c_{n+2} - a \sum_{j=0}^n (n+1-j) c_j c_{n+1-j} \right). \end{aligned} \quad (19)$$

Thus, the power series solution of (16) is as follows

$$\begin{aligned} \Psi(\theta) = & c_0 + c_1 \theta + c_2 \theta^2 + c_3 \theta^3 + \sum_{n=1}^{\infty} c_{n+3} \theta^{n+3} \\ = & c_0 + c_1 \theta + c_2 \theta^2 + \frac{\lambda c_1 - ac_0 c_1 - 2(d+e)c_2}{6(b+c)} \theta^3 \\ & + \sum_{n=1}^{\infty} \frac{1}{(b+c)(n+1)(n+2)(n+3)} \left(\lambda(n+1)c_{n+1} \right. \\ & \left. - (d+e)(n+1)(n+2)c_{n+2} - a \sum_{j=0}^n (n+1-j) c_j c_{n+1-j} \right) \theta^{n+3}. \end{aligned} \quad (20)$$

Consequently, the exact power series solution of (1) can be written as follows

$$u(x, y, t) = C_0 + C_1(x - y - \lambda t) + C_2(x - y - \lambda t)^2 + C_3(x - y - \lambda t)^3 + \sum_{n=1}^{\infty} C_{n+3}(x - y - \lambda t)^{n+3} = C_0 + C_1(x - y - \lambda t) + C_2(x - y - \lambda t)^2 + \frac{\lambda C_1 - a C_0 C_1 - 2(d + e)C_2}{6(b + c)}(x - y - \lambda t)^3 + \sum_{n=1}^{\infty} \frac{1}{(b + c)(n + 1)(n + 2)(n + 3)}(\lambda(n + 1)C_{n+1} - (d + e)(n + 1)(n + 2)C_{n+2} - a \sum_{j=0}^n (n + 1 - j)C_j C_{n+1-j})(x - y - \lambda t)^{n+3}, \quad (21)$$

where $C_i (i = 0, 1, 2, 3)$ are arbitrary constants, the other coefficients $C_n (n \geq 3)$ also can be derived.

Remark 1. The explicit solutions of other equations can also be derived in the same way. Here we do not list them for simplicity.

In order to search for others explicit solutions of Eq. (1), we make use of the auxiliary equation, i.e., the Riccati equation of the following "general form"

$$\varphi' = r + p\varphi + q\varphi^2, \quad (22)$$

and use its solution to construct the solutions for ZKB equation. Here, p, q, r are real constant. By omitting the details, we directly write the general solution of (22) as

$$\varphi = \frac{\sqrt{4rq - p^2}}{2q} \frac{C_1 e^{\frac{\theta}{2}\sqrt{4rq - p^2}} - C_2 e^{-\frac{\theta}{2}\sqrt{4rq - p^2}}}{C_1 e^{\frac{\theta}{2}\sqrt{4rq - p^2}} + C_2 e^{-\frac{\theta}{2}\sqrt{4rq - p^2}}} - \frac{p}{2q}, \quad (23)$$

where C_1, C_2 are arbitrary constants. By balancing the highest derivative and nonlinear terms in (15), we assume the solution of (1) of the form

$$f(\theta) = a_0 + a_1\varphi + a_2\varphi^2, \quad (24)$$

where a_0, a_1, a_2 are constants to be determined.

Substituting the ansatz (24) along with (22) into (15), collecting coefficients of monomials of φ with the aid of Maple, and then setting each coefficients equal to zero, one gets

$$a_2 = -12 \frac{q^2(b + c)}{a}, \quad a_1 = -\frac{12}{5} \frac{(5bp + 5cp - d - e)q}{a}, \quad \lambda = \frac{5aa_0 - 6p(d + e) + 60qr(b + c)}{5}, \quad (25)$$

where a_0 is a arbitrary constant. Also,

$$(d + e)^2 = 25p^2(b + c)^2 + 100qr(b + c)^2. \quad (26)$$

From the ansatz (24) and making use of Eqs. (25) and (23), one can get the explicit solution of (1)

$$u(x, y, t) = -12 \frac{q^2(b + c)}{a} \left(\frac{\sqrt{4rq - p^2}}{2q} \frac{C_1 e^{\frac{\theta}{2}\sqrt{4rq - p^2}} - C_2 e^{-\frac{\theta}{2}\sqrt{4rq - p^2}}}{C_1 e^{\frac{\theta}{2}\sqrt{4rq - p^2}} + C_2 e^{-\frac{\theta}{2}\sqrt{4rq - p^2}}} - \frac{p}{2q} \right)^2 - \frac{12}{5} \frac{(5bp + 5cp - d - e)q}{a} \left(\frac{\sqrt{4rq - p^2}}{2q} \frac{C_1 e^{\frac{\theta}{2}\sqrt{4rq - p^2}} - C_2 e^{-\frac{\theta}{2}\sqrt{4rq - p^2}}}{C_1 e^{\frac{\theta}{2}\sqrt{4rq - p^2}} + C_2 e^{-\frac{\theta}{2}\sqrt{4rq - p^2}}} - \frac{p}{2q} \right) + a_0, \quad (27)$$

here $\theta = x - y - \lambda t$, and λ is given by (25).

The solution (27) is the general solution of ZKB equation and therefore several independently real solutions can be obtained. For instance, the exact solution (21) which was obtained by "tanh method" in Ref. [27], can be retrieved from (27) by choosing $C_1 = C_2 = 1, p = 0, a_0 = \frac{9(d + e)^2}{25a(b + c)}$, and $qr = \frac{(d + e)^2}{100(b + c)^2}$. Similarly, one can compare the accuracy of the numerical results obtained in [26] via (27).

In particular, if we let $C_1 = 1, C_2 = -1, p = 0$, and $4rq - p^2 > 0$, one can get

$$u(x, y, t) = -12 \frac{q^2(b + c)}{a} \left(\frac{\sqrt{4rq - p^2}}{2q} \cot h \left(\frac{\theta}{2} \sqrt{4rq - p^2} \right) - \frac{p}{2q} \right)^2 - \frac{12}{5} \frac{(5bp + 5cp - d - e)q}{a} \left(\cot h \left(\frac{\theta}{2} \sqrt{4rq - p^2} \right) - \frac{p}{2q} \right) + a_0. \quad (28)$$

In particular, if we set $C_1 = 1, C_2 = 1$, and $4rq - p^2 > 0$, one can derive

$$u(x, y, t) = -12 \frac{q^2(b + c)}{a} \left(\frac{\sqrt{4rq - p^2}}{2q} \tan h \left(\frac{\theta}{2} \sqrt{4rq - p^2} \right) - \frac{p}{2q} \right)^2 - \frac{12}{5} \frac{(5bp + 5cp - d - e)q}{a} \left(\tan h \left(\frac{\theta}{2} \sqrt{4rq - p^2} \right) - \frac{p}{2q} \right) + a_0. \quad (29)$$

3. Conservation laws of (1)

In this section, we obtain conservation laws for the ZKB equation.

Theorem 1 ([30,31]). The system and its adjoint equation

$$F(x, u, u_{(1)}, u_{(2)}, \dots, u_{(s)}) = 0, \quad F^*(x, u, v, u_{(1)}, v_{(1)}, u_{(2)}, v_{(2)}, \dots, u_{(s)}, v_{(s)}) = 0, \quad (29)$$

has a formal Lagrangian, namely

$$L = vF(x, u, u_{(1)}, u_{(2)}, \dots, u_{(s)}). \quad (30)$$

In the following we recall the "new conservation theorem" given by Ibragimov in [31].

Definition 1. [30,31] The first equation of (29) is said to be nonlinearly self-adjoint if for some arbitrary function $\phi(x, u) \neq 0$, we have

$$F^*|_{v=\phi} = \lambda(x, u, u_{(1)}, \dots)F, \quad (31)$$

where λ is an indeterminate variable coefficient.

Theorem 2 [32]. Every Lie point, Lie-Bäcklund and non-local symmetry of Eq. (1) provides a conservation law for Eq. (1) and the adjoint equation. Then the elements of conservation vector (C^1, C^2, C^3) are given by

$$C^i = \zeta^i L + W^\alpha \left[\frac{\partial L}{\partial u_i^\alpha} - D_j \left(\frac{\partial L}{\partial u_{ij}^\alpha} \right) + D_j D_k \left(\frac{\partial L}{\partial u_{ijk}^\alpha} \right) \right] + D_j (W^\alpha) \left[\left(\frac{\partial L}{\partial u_{ij}^\alpha} \right) - D_k \left(\frac{\partial L}{\partial u_{ijk}^\alpha} \right) + \dots \right], \quad (32)$$

where $W^\alpha = \eta^\alpha - \zeta^j u_{j\alpha}^\alpha$.

3.1. Nonlinear self-adjointness

For (1), the adjoint equation has the form

$$F^* = -(v_t + auv_x + bv_{xxx} + cv_{xyy} + dv_{xx} + ev_{yy}) = 0, \quad (33)$$

and the Lagrangian in the symmetrized form

$$L = v(u_t + auu_x + bu_{xxx} + cu_{xyy} - du_{xx} - eu_{yy}). \quad (34)$$

If we substitute u instead of v in Eq. (33), one can find that Eq. (1) is not recovered. Therefore, we can say Eq. (1) is not self adjoint. Next,

we look for an explicit form of $\phi(x, u) \neq 0$ for Eq. (1) that holds Eq. (33).

$$v_t + auv_x + bv_{xxx} + cv_{xyy} + dv_{xx} + ev_{yy}|_{v=\phi(x,u)} = \lambda(u_t + auu_x + bu_{xxx} + cu_{xyy} - du_{xx} - eu_{yy}). \quad (35)$$

If we set $v = \phi(t, x, y, u)$, one can get

$$\begin{aligned} v_t &= \phi_t + \phi_u u_t, \\ v_x &= \phi_x + \phi_u u_x, \\ v_{xx} &= \phi_{xx} + 2\phi_{xu} u_x + \phi_{uu} u_x^2 + \phi_u u_{xx}, \\ v_{yy} &= \phi_{yy} + 2\phi_{yu} u_y + \phi_{uu} u_y^2 + \phi_u u_{yy}, \\ v_{xxx} &= \phi_{xxx} + 3\phi_{xxu} u_x + 3\phi_{xuu} u_x^2 + 3\phi_{xu} u_{xx} + \phi_{uuu} u_x^3 \\ &\quad + 3\phi_{uu} u_x u_{xx} + \phi_u u_{xxx}, \\ v_{xyy} &= \phi_{xyy} + \phi_{yyu} u_x + 2\phi_{xyu} u_y + 2\phi_{yuu} u_x u_y + 2\phi_{yu} u_{xy} \\ &\quad + \phi_{uux} u_y^2 + \phi_{uuu} u_x u_y^2 + 2\phi_{uu} u_y u_{yx} + \phi_{ux} u_{yy} \\ &\quad + \phi_{uu} u_x u_{yy} + \phi_u u_{xyy}. \end{aligned} \quad (36)$$

Plugging them into (35), one can arrive at the following self-adjointness condition

$$\begin{aligned} &\phi_t + \phi_u u_t + au(\phi_x + \phi_u u_x) + d(\phi_{xx} + 2\phi_{xu} u_x + \phi_{uu} u_x^2 + \phi_u u_{xx}) \\ &\quad + e(\phi_{yy} + 2\phi_{yu} u_y + \phi_{uu} u_y^2 + \phi_u u_{yy}) + b(\phi_{xxx} + 3\phi_{xxu} u_x + 3\phi_{xuu} u_x^2 \\ &\quad + 3\phi_{xu} u_{xx} + \phi_{uuu} u_x^3 + 3\phi_{uu} u_x u_{xx} + \phi_u u_{xxx}) + c(\phi_{xyy} + \phi_{yyu} u_x + 2\phi_{xyu} u_y \\ &\quad + 2\phi_{yuu} u_x u_y + 2\phi_{yu} u_{xy} + \phi_{uux} u_y^2 + \phi_{uuu} u_x u_y^2 + 2\phi_{uu} u_y u_{yx} \\ &\quad + \phi_{ux} u_{yy} + \phi_{uu} u_x u_{yy} + \phi_u u_{xyy}) = \lambda(u_t + auu_x + bu_{xxx} + cu_{xyy} - du_{xx} - eu_{yy}). \end{aligned} \quad (37)$$

From the coefficient of the term u_t , one can get

$$\lambda = -\phi_u. \quad (38)$$

Note that the other coefficients of u and all of the derivative yields

$$\begin{aligned} b\phi_{uuu} &= 0, \quad c\phi_{uuu} = 0, \quad 3b\phi_{uu} = 0, \quad 2c\phi_{yuu} = 0, \quad \phi_{yu} = 0, \\ 3b\phi_{xxu} + 2d\phi_{xu} + c\phi_{yyu} &= 0, \quad 2ec\phi_{yu} + 2c\phi_{xyu} = 0, \\ 3b\phi_{xu} + 2d\phi_u &= 0, \quad c\phi_{xu} + 2e\phi_u = 0, \\ 3b\phi_{xuu} + d\phi_{uu} &= 0, \quad c\phi_{xuu} + e\phi_{uu} = 0, \\ \phi_t + au\phi_x + b\phi_{xxx} + d\phi_{xx} + e\phi_{yy} + c\phi_{xyy} &= 0. \end{aligned} \quad (39)$$

Solve them, one can get the solution

$$\phi = \frac{c_3 \left((e^{\sqrt{k_1} y})^2 c_1 + c_2 \right)}{e^{\sqrt{k_1} y} e^{k_1 e t}}. \quad (40)$$

In particular, (39) has a special solution

$$\phi = c_1 y + c_2, \quad (41)$$

where c_1, c_2 and c_3 are constants. We, thus, have the following statement.

Theorem 3. The ZKB Eq. (1) is nonlinearly self-adjoint with the substitution $v = \phi$ and ϕ given by (40) or (41).

3.2. Conservation laws of (1)

We now construct the conservation laws by using the adjoint equation and symmetries of (1). For (1), the adjoint equation is given by

$$F = v_t + auv_x + bv_{xxx} + cv_{xyy} + dv_{xx} + ev_{yy}, \quad (42)$$

and the Lagrangian in the symmetrized form

$$L = v \left(u_t + auu_x + bu_{xxx} + \frac{1}{3} c(u_{xyy} + u_{yxy} + u_{yyx}) - du_{xx} - eu_{yy} \right). \quad (43)$$

Consider Theorem 2, the corresponding vector fields is given by

$$\begin{aligned} V &= \xi^1(x, y, t, u) \frac{\partial}{\partial t} + \xi^2(x, y, t, u) \frac{\partial}{\partial x} + \xi^3(x, y, t, u) \frac{\partial}{\partial y} \\ &\quad + \phi(x, y, t, u) \frac{\partial}{\partial u}. \end{aligned} \quad (44)$$

The conservation law is decided by

$$D_t(C^1) + D_x(C^2) + D_y(C^3) = 0. \quad (45)$$

Here the conserved vector $C = (C^1, C^2, C^3)$ are given by (32) and the components given by

$$C^1 = \xi^1 L + W \frac{\partial L}{\partial u_t}, \quad (46)$$

$$\begin{aligned} C^2 &= \xi^2 L + W \left[\frac{\partial L}{\partial u_x} - D_x \frac{\partial L}{\partial u_{xx}} + D_{xx} \frac{\partial L}{\partial u_{xxx}} + D_{yy} \frac{\partial L}{\partial u_{xyy}} \right] \\ &\quad + W_x \left[\frac{\partial L}{\partial u_{xx}} - D_x \frac{\partial L}{\partial u_{xxx}} \right] + W_y \left[-D_y \frac{\partial L}{\partial u_{xyy}} \right] + W_{xx} \left[\frac{\partial L}{\partial u_{xxx}} \right] \\ &\quad + W_{yy} \left[\frac{\partial L}{\partial u_{xyy}} \right], \end{aligned} \quad (47)$$

$$\begin{aligned} C^3 &= \xi^3 L + W \left[-D_y \frac{\partial L}{\partial u_{yy}} + D_{xy} \frac{\partial L}{\partial u_{yxy}} + D_{yx} \frac{\partial L}{\partial u_{yyx}} \right] \\ &\quad + W_x \left[-D_y \frac{\partial L}{\partial u_{yxy}} \right] + W_y \left[\frac{\partial L}{\partial u_{yy}} - D_x \frac{\partial L}{\partial u_{yyx}} \right] + W_{xy} \left[\frac{\partial L}{\partial u_{yyx}} \right] \\ &\quad + W_{yx} \left[\frac{\partial L}{\partial u_{yyx}} \right], \end{aligned} \quad (48)$$

that is

$$C^1 = \xi^1 L + W v, \quad (49)$$

$$\begin{aligned} C^2 &= \xi^2 L + W \left[auv - D_x(-dv) + D_{xx}(bv) + D_{yy} \left(\frac{1}{3} cv \right) \right] \\ &\quad + W_x(-dv - D_x(bv)) + W_y \left(-D_y \left(\frac{1}{3} cv \right) \right) + W_{xx}bv \\ &\quad + W_{yy} \frac{1}{3} cv \\ &= \xi^2 L + W \left(auv + dv_x + bv_{xx} + \frac{1}{3} cv_{yy} \right) - W_x(dv + bv_x) \\ &\quad - W_y \left(\frac{1}{3} cv_y \right) + bvW_{xx} + \frac{1}{3} cvW_{yy}, \end{aligned} \quad (50)$$

$$\begin{aligned} C^3 &= \xi^3 L + W \left[-D_y(-ev) + D_{xy} \left(\frac{2}{3} cv \right) \right] + W_x \left[-D_y \left(\frac{1}{3} cv \right) \right] \\ &\quad + W_y \left[-ev - D_x \left(\frac{1}{3} cv \right) \right] + W_{xy} \left[\frac{2}{3} cv \right] \\ &= \xi^3 L + W \left(ev_y + \frac{2}{3} cv_{xy} \right) - \frac{1}{3} cv_y W_x - W_y \left(ev + \frac{1}{3} cv_x \right) \\ &\quad + \frac{2}{3} cvW_{xy}, \end{aligned} \quad (51)$$

with

$$W = \phi - \xi^1 u_t - \xi^2 u_x - \xi^3 u_y. \quad (52)$$

Next, we consider following cases.

Case 1.

For the operator $V = \frac{\partial}{\partial t}$, we have

$$W = -u_t, \quad (53)$$

we can get the conservation vector of (1)

$$C^1 = -u_t v, \quad (54)$$

$$C^2 = -u_t \left(auv + dv_x + bv_{xx} + \frac{1}{3} cv_{yy} \right) + u_{tx} (dv + bv_x) + \frac{1}{3} cu_{ty} v_y - bvu_{txx} - \frac{1}{3} cvu_{tyy}, \quad (55)$$

$$C^3 = -u_t \left(ev_y + \frac{2}{3} cv_{xy} \right) + \frac{1}{3} cv_y u_{tx} + u_{ty} \left(ev + \frac{1}{3} cv_x \right) - \frac{2}{3} cvu_{txy}. \quad (56)$$

Case 2.

For the generator $V = \frac{\partial}{\partial x}$, we get

$$W = -u_x, \quad (57)$$

one can obtain the conservation vector of (1)

$$C^1 = -u_x v, \quad (58)$$

$$C^2 = -u_x \left(auv + dv_x + bv_{xx} + \frac{1}{3} cv_{yy} \right) + u_{xx} (dv + bv_x) + \frac{1}{3} cu_{xy} v_y - bvu_{xxx} - \frac{1}{3} cvu_{xyy}, \quad (59)$$

$$C^3 = -u_x \left(ev_y + \frac{2}{3} cv_{xy} \right) + \frac{1}{3} cv_y u_{xx} + u_{xy} \left(ev + \frac{1}{3} cv_x \right) - \frac{2}{3} cvu_{xxy}. \quad (60)$$

Case 3.

For the Lie algebra $V = \frac{\partial}{\partial y}$, one can arrive at

$$W = -u_y, \quad (61)$$

we can reach the conservation vector of (1)

$$C^1 = -u_y v, \quad (62)$$

$$C^2 = -u_y \left(auv + dv_x + bv_{xx} + \frac{1}{3} cv_{yy} \right) + u_{yx} (dv + bv_x) + \frac{1}{3} cu_{yy} v_y - bvu_{yxx} - \frac{1}{3} cvu_{yyy}, \quad (63)$$

$$C^3 = -u_y \left(ev_y + \frac{2}{3} cv_{xy} \right) + \frac{1}{3} cv_y u_{yx} + u_{yy} \left(ev + \frac{1}{3} cv_x \right) - \frac{2}{3} cvu_{xyy}. \quad (64)$$

Case 4.

For the operator $V = t \frac{\partial}{\partial x} + \frac{1}{a} \frac{\partial}{\partial u}$, we have

$$W = \frac{1}{a} - tu_x, \quad (65)$$

we derive the conservation vector of (1)

$$C^1 = \frac{1}{a} v - tu_x v, \quad (66)$$

$$C^2 = \left(\frac{1}{a} - tu_x \right) \left(auv + dv_x + bv_{xx} + \frac{1}{3} cv_{yy} \right) + tu_{xx} (dv + bv_x) + \frac{1}{3} ctu_{xy} v_y - btvu_{xxx} - \frac{1}{3} ctvu_{xyy}, \quad (67)$$

$$C^3 = \left(\frac{1}{a} - tu_x \right) \left(ev_y + \frac{2}{3} cv_{xy} \right) + \frac{1}{3} ctv_y u_{xx} + tu_{xy} \left(ev + \frac{1}{3} cv_x \right) - \frac{2}{3} ctvu_{xyx}. \quad (68)$$

It is clear that they are involves an arbitrary solution v of the adjoint Eq. (33), and they presents an infinite number of the conservation laws.

4. Conclusions

In the present paper, by using the Lie symmetry groups, we studied the symmetry properties, similarity reduction forms and explicit solutions of the (2+1)-dimensional ZKB equation. Moreover, we also constructed the power series solution of the equation. At last, we derived the nonlinear self-adjointness of Eq. (1), by virtue of this fact, infinitely many conservation laws of the equation are exhibited. Furthermore, the results obtained here can be useful in enhancing the understanding of nonlinear propagation of small amplitude electrostatic structures in dense, dissipative (e-p-i) magnetoplasmas.

Acknowledgement

We are extremely thankful for the reviewers for their many valuable suggestions that greatly improved this paper.

References

- [1] Olver PJ. Application of lie group to differential equation. New York: Springer; 1986.
- [2] Ovsiannikov LV. Group analysis of differential equations. New York: Academic Press; 1982.
- [3] Bluman GW, Kumei S. Symmetries and differential equations. New York: Springer; 1989.
- [4] Li YS, Ma WX, Zhang JE. Darboux transformations of classical Boussinesq system and its new solutions. Phys Lett A 2000;275:60–6.
- [5] Zayed EME, Zedan HA, Gepreel KA. On the solitary wave solutions for nonlinear Hirota–Satsuma coupled KdV of equations. Chaos, Solitons Fractals 2004;22:285–303.
- [6] Kumara S, Singhb K, Gupta RK. Painleve analysis, Lie symmetries and exact solutions for (2+1)-dimensional variable coefficients Broer–Kaup equations. Commun Nonlinear Sci Numer Simulat 2012;17:1529–41.
- [7] Ablowitz MJ, Segur H. Solitons and inverse scattering transform. Philadelphia: SIAM; 1981.
- [8] Fuchssteiner Benno, Fokas Athanassios S. Symplectic structures, their Backlund transformations and hereditary symmetries. Phys D: Nonlinear Phenom 1981;4:47–66.
- [9] Avdonina ED, Ibragimov NH, Khamitova R. Exact solutions of gasdynamic equations obtained by the method of conservation laws. Commun Nonlinear Sci Numer Simulat 2013;18:2359–66.
- [10] Hirota R. The DirectMethod in soliton theory. Cambridge: Cambridge University Press; 2004.
- [11] Biswas A, Kara AH, Bokhari AH, Zaman FD. Solitons and conservation laws of Klein–Gordon equation with power law and log law nonlinearities. Nonlinear Dyn 2013;73:2191–6.
- [12] Zakharov VE, Kuznetsov EA. On three-dimensional solitons. Sov Phys -JETP 1974;39:285–8 [Appl Math Comput].
- [13] Biswas A. 1-Soliton solution of the generalized Zakharov–Kuznetsov modified equal width equation. Appl Math Lett 2009;22:1775–7.
- [14] Biswas A, Zerrad E. 1-Soliton solution of the Zakharov–Kuznetsov equation with dual-power law nonlinearity. Commun Nonlinear Sci Numer Simulat 2009;14:3574–7.
- [15] Biswas A. 1-Soliton solution of the generalized Zakharov–Kuznetsov equation with nonlinear dispersion and time-dependent coefficients. Phys Lett A 2009;373:2931–4.
- [16] Krishnan EV, Biswas A. Solutions to the Zakharov–Kuznetsov equation with higher order nonlinearity by mapping and ansatz methods. Phys Wave Phenom 2010;18:256–61.
- [17] Johnpillai AG, Kara AH, Biswas A. Symmetry solutions and reductions of a class of generalized (2+1)-dimensional Zakharov–Kuznetsov equation. Int J Nonlin Sci Num 2011;12:35–43.
- [18] Ebadi G, Mojaver A, Milovic Da, Johnson S, Biswas A. Solitons and other solutions to quantum Zakharov–Kuznetsov equation in quantum magnetoplasmas. Astrophys Space Sci 2011;341:507–13.

- [19] Bhrawy AH, Abdelkawy MA, Kumar S, Johnson S, Biswas A. Solitons and other solutions to quantum Zakharov–Kuznetsov equation in quantum magnetoplasmas. *Indian J Phys* 2013;87:455–63.
- [20] Stenflo L, Shukla PK, Marklund M. New low-frequency oscillations in quantum dusty plasmas. *Europhys Lett* 2006;74:844–6.
- [21] Khan SA, Masood W, Siddiq M. Obliquely propagating dust-acoustic waves in dense quantum magnetoplasmas. *Phys Plasmas* 2009;16:013701.
- [22] Hasegawa H, Irie S, Usami S, Ohsawa Y. Perpendicular nonlinear waves in an electronpositronion plasma. *Phys Plasmas* 2002;9:2549–61.
- [23] Masood W, Rizvi H, Hussain S. Two dimensional electromagnetic shock structures in dense electron-positron-ion magnetoplasmas. *Astrophys Space Sci* 2011;332:287–99.
- [24] Mamun AA, Shukla PK. Cylindrical and spherical dust ion-coustic solitary waves. *Phys Plasmas* 2002;9:1468–70.
- [25] Shukla PK, Mamun AA. Solitons, shocks and vortices in dusty plasmas. *New J Phys* 2003;5:17.
- [26] El-Bedwehy NA, Moslem WM. Zakharov–Kuznetsov–Burgers equation in superthermal electron-positron-ion plasma. *Astrophys Space Sci* 2011;335:435–42.
- [27] Masood W, Rizvi H, Siddiq M. Obliquely propagating nonlinear structures in dense dissipative electron positron ion magnetoplasmas. *Astrophys Space Sci* 2012;337:629–35.
- [28] Liu H, Li J. Lie symmetry analysis and exact solutions for the short pulse equation. *Nonlinear Anal* 2009;71:2126–33.
- [29] Wang GW, Liu XQ, Zhang YY. Symmetry reduction, exact solutions and conservation laws of a new fifth-order nonlinear integrable equation. *Commun Nonlinear Sci Numer Simulat* 2013;18:2313–20.
- [30] Ibragimov NH. Integrating factors, adjoint equations and Lagrangians. *J Math Anal Appl* 2006;318:742–57.
- [31] Ibragimov NH. Nonlinear self-adjointness and conservation laws. *J Phys A: Math Theor* 2011;44:432002.
- [32] Ibragimov NH. A new conservation theorem. *J Math Anal Appl* 2007;333:311–28.

524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541